

3.0 RESULTS AND DISCUSSION

3.1 STRATIFICATION MONITORING

The following sections summarize the water quality monitoring results of the physical and chemical parameters: temperature, dissolved oxygen, pH, and conductivity. For each parameter, we describe seasonal and spatial patterns of surface water quality measured throughout the reservoir watershed, and seasonal and depth related patterns of the stratified water column based on measures from the deepest portion of the reservoir (station BZ-6 or the "Tower"). We feel that it is appropriate to focus discussion of stratification at this station as water quality problems related to depth are generally most severe in deeper water habitats, thus our evaluation will be a conservative one. Finally, we analyze 2001 data along with the Beltzville Reservoir historical database for trends in dissolved oxygen concentrations over the past two decades. Versar personnel collected the physical and chemical water quality data discussed herein over the monitoring period from April to October 2001. All of the parameters were measured with a calibrated Hydrolab water quality meter and are presented in Appendix Table A.

3.1.1 Temperature

Temperatures of surface water generally followed a similar pattern throughout the watershed of Beltzville Reservoir during 2001, however consistent differences were apparent between reservoir-body and upstream and downstream waters (Fig. 3-1). Stations located in the reservoir-body (BZ-3, -6, and -7) averaged 20 °C throughout the monitoring period. Temperatures downstream of the reservoir (BZ-1) and at the upstream stations (BZ-2, -4, -5) were similar but averaged 6 °C cooler than the reservoir.

Beltzville Reservoir was stratified with respect to temperature in 2001 (Fig. 3-2). In April stratification was already apparent with surface temperatures (10 °C) approximately 4 °C warmer than the lower water column (6 °C). This difference was also observed in May, however the upper water column from the surface to about 50 feet had warmed to about 15 °C. Throughout June and into September, the temperature pattern was very consistent with temperature at the surface averaging 22 °C and decreasing to 6 °C near the bottom.

3.1.2 Dissolved Oxygen

Dissolved oxygen (DO) in the surface water generally followed a similar pattern throughout the watershed of Beltzville Reservoir during 2001, however consistent differences were apparent between reservoir-body and upstream and downstream waters (Fig. 3-3). Stations located in the reservoir-body (BZ-3, -6, and -7) averaged 8-mg/L throughout the monitoring period. Dissolved Oxygen concentrations downstream of the reservoir




Figure 3-1. Surface water temperature (°C) measured at Beltzville Reservoir in 2001. See Appendix A for Summary of plotted values.

Temperature - Stratification

3-3




Figure 3-2. Temperature stratification at station BZ-6 of Beltzville Reservoir in 2001. See Appendix A for summary of plotted values.

Dissolved Oxygen - Surface Water




Figure 3-3. Dissolved oxygen concentrations measured in surface waters at Beltzville Reservoir in 2001. (The PADEP water quality standard for dissolved oxygen is a minimum concentration of 5 mg/L.) See Appendix A for summary of plotted values.

(BZ-1) and at the upstream stations (BZ-2, -4, -5) were consistently higher and averaged 9-mg/L.

DO in the water column of Beltzville Reservoir was stratified during September and October (Fig. 3-4). However, in most months DO concentrations were high throughout the water column. In September and October, DO concentrations were severely depleted throughout most of the water column. In these months, DO was less than 5-mg/L from 45-ft to the bottom, while concentrations nearer the surface were closer to 8-mg/L. Also during these months, DO exhibited a sinuous pattern with concentrations decreasing rapidly from the surface to approximately 50 feet, increasing to a depth of 90 feet, and decreasing again thereafter to the bottom. This pattern has been observed in previous years and may be due to a lens of oxygenated water passing through the reservoir from upstream sources or a result of portal operations at the reservoir tower.

DO concentrations in the water column of Beltzville Reservoir at times were not in compliance with PADEP water quality standards during 2001. The state water quality standard for DO is a minimum concentration of 5-mg/L. As stated above, concentrations falling below the standard were encountered during the September and October. In addition, the concentrations of DO fell to 4.6-mg/L during August (Fig 3-4). DO concentrations measured in all surface waters of the reservoir and upstream and downstream were in compliance with the standard.

The health of aquatic ecosystems is impaired by low DO concentrations in the water column. Hypoxia, or conditions of DO less than 2 mg/L, is generally accepted as the threshold at which the most severe effects on biota occur. In 2001, the deeper water column of Beltzville was affected by hypoxia (Fig. 3-5). Hypoxic water was encountered in September and October and occupied the lower two-thirds of the water column from around 45-ft to the bottom. Hypoxia in the lower water column is a symptom of eutrophication. Nutrients in the water column feed accelerating algal growth at the surface photic zone. Dead and decaying algae sink to lower levels of the water column and during the process of decay; oxygen is depleted from the water column. This phenomenon is not uncommon for lake systems.

Since 1988, DO concentrations have been measured in the water column at station BZ-6 near the "Tower". Station BZ-6 is located in the deepest part of the reservoir (120-ft) at which DO has been measured at 5-foot intervals from the surface to the bottom. We analyzed this historical data along with this year's data to determine if the depth at which hypoxia was initially encountered in the water column had significantly changed over the past 14 years. The trend analysis was conducted for the summer season (July to September) only as hypoxia was rarely observed in the spring (April to June). From the regression analysis, the level of hypoxia has not changed significantly ($P > 0.05$; $R^2 = 0.00003$) over the past 14 years (Fig. 3-6). In most years, the average depth at which hypoxia occurred was between 70 and 90 feet. In 2001, this average depth at which hypoxia occurred was 80 feet.

Dissolved Oxygen - Stratification

3-6





Figure 3-4. Dissolved oxygen stratification at station BZ-6 of Beltzville Reservoir in 2001. (The PADEP water quality standard for DO is a minimum concentration of 5 mg/L.) See Appendix A for summary of plotted values.

Tower - Hypoxia



3-7

Figure 3-5. Spatial/temporal distribution of hypoxic reservoir water in Beltzville Reservoir measured at the "Tower" station in 2001. Histograms indicate dissolved oxygen concentrations in the water column below 2.0 mg/L.

Tower - Hypoxia Trends




Figure 3-6. Trends in seasonal average hypoxia at the "Tower" station BZ-6. Measured as the depth at which hypoxia was observed in the water column.

In addition to the regression analysis, we also used the Mann-Kendall test to determine if DO concentrations had changed significantly at individual monitoring stations over the past 20 years or more. As before, the trend analysis was conducted separately for spring (April to June) and summer (July to September) seasons. Surface water concentrations were analyzed at upstream station BZ-5 and downstream station BZ-1. Surface and bottom water concentrations were averaged at the reservoir-body station BZ-3. Dissolved oxygen in surface waters of Beltzville Reservoir does not appear to have changed in concentration over the past two decades. However, a significant decreasing trend was determined from the analysis at station BZ-3 for the summer. The rate of decrease was 0.12 mg/L/year (Table 3-1).

Table 3-1. Spring and summer season trends of dissolved oxygen concentration at individual stations of Beltzville Reservoir calculated with the Mann-Kendall Statistic.					
Station	# of Years spring/summer	Spring		Summer	
		P Level	Rate (mg/L/year)	P Level	Rate (mg/L/year)
Surface Water					
BZ-1	23	NS	0.02	NS	-0.02
BZ-2	21	NS	0.03	NS	0.01
BZ-3	25	NS	-0.02	< 0.05	-0.12
BZ-4	23	NS	0.01	NS	0.01
BZ-5	24/25	NS	0.02	NS	0.01
BZ-6	6	NS	-0.24	NS	0.22
BZ-7	6	NS	-0.12	NS	0.17

3.1.3 pH

The concentrations of pH in surface water followed a similar pattern at Beltzville Reservoir during 2001 (Fig. 3-7). Throughout the reservoir and upstream and downstream stations the pH among stations remained consistent among stations. The overall average pH throughout the monitoring period was 7.1, with a maximum average of 7.3 at station BZ-7.

In 2001, pH in the water column of Beltzville Reservoir was broadly effected by stratification (Fig. 3-8). In general, the stratification pattern was marked by a rapidly decreasing pH from the surface to about 30 feet and thereafter remaining relatively consistent to the bottom. In April, pH at the surface was 7.2 while the lower water column was about 6.9. In early August, pH of the entire water column was lowest and ranged from 7.5 at the surface to about 5.6 in the lower water column.

pH - Surface Water




Figure 3-7. Surface water pH measured at Beltzville Reservoir in 2001. (The PADEP water quality standard for pH is from 6 to 9.) See Appendix A for summary of plotted values.

pH - Stratification




Figure 3-8. pH stratification at station BZ-6 of Beltzville Reservoir in 2001. (The PADEP water quality standard for pH is from 6 to 9.) See Appendix A for summary of plotted values.

The pH measures in the lower water column of Beltzville Reservoir were not in compliance with PADEP water quality standards from July to September. The standard for pH is a range of acceptable measures between 6 and 9. In July, August, and September, pH at station BZ-6 were below the minimum value of the standard. In those months, the average pH at station BZ-6 below 30 feet was 5.9. Additionally, the average pH at station BZ-7 below 25 feet in August was 5.9.

3.1.4 Conductivity

Conductivity measured in the surface waters of Beltzville Reservoir was generally consistent during 2001. Measures in the reservoir-body (stations BZ-3, BZ-6, and BZ-7), downstream of the reservoir (BZ-1), and upstream stations (BZ-2 and BZ-5) were most similar and averaged about 0.08-mS/cm throughout the monitoring period (Fig. 3-9). Conductivity measured at Wild Creek (BZ-4) was consistently the lowest, averaging 0.04-mS/cm and ranged from 0.07-mS/cm in May to 0.09-mS/cm in September.

Conductivity throughout the water column of Beltzville Reservoir was also fairly consistent during 2001 (Fig. 3-10). In May, measures were most consistent at about 0.074-mS/cm from surface to bottom. In the later months of August, September, and October, measures were lower at the surface than the bottom and overall ranged from 0.070-mS/cm to 0.089-mS/cm. The increasing conductivity in the lower water column during the summer months is an indication of eutrophic influences. As oxygen is depleted in the lower water column, a chemically reducing environment is formed. In this state, metals are mobilized into the water column, which results in an increase in conductivity. Conversely, at the surface, biological activity in the photic zone during the summer months effectively reduces conductivity as metabolizing algae remove dissolved constituents.

3.2 WATER COLUMN CHEMISTRY MONITORING

The following sections describe temporal, spatial, and patterns relating to depth for the water quality parameters measured in surface, middle, and bottom waters of Beltzville Reservoir during 2001 (Table 3-2). Where appropriate, long-term trends are discussed for surface water quality parameters incorporating 2001 data with the Beltzville Reservoir historical water quality database.

3.2.1 Ammonia

Ammonia concentrations were generally low in Beltzville Reservoir during 2001 (Fig. 3-11). Concentrations at most stations and depths were near or less than the method detection limit of 0.1 mg/L. The highest concentrations of ammonia (0.3-mg/L) were measured in the surface water at station BZ-1 during April and the middle water at station BZ-6 during August. Elevated ammonia in the lower water column of deep, stratified lakes and

Conductivity - Surface




Figure 3-9. Surface water conductivity (mS/cm) measured at Beltzville Reservoir in 2001. See Appendix A for summary of plotted values.

Conductivity - Stratification

3-14




Figure 3-10. Conductivity stratification at station BZ-6 of Beltzville Reservoir in 2001. See Appendix A for summary of plotted values.

Table 3-2. Summary of surface, middle, and bottom water quality monitoring data for Beltzville Reservoir in 2001													
STATION	DATE	NH3	NO2	NO3	PO4	TKN	TP	TDS	TSS	BOD5	ALK	CHL A	
BZ-1S	24-Apr	0.30	< 0.005	0.100	< 0.05	0.30	< 0.05	28	8.0	< 3.0	10.0	2.3	
	24-May	< 0.10	< 0.005	0.700	< 0.05	< 0.10	< 0.05	96	2.0	< 3.0	8.0	1.5	
	14-Jun	< 0.10	< 0.005	< 0.100	< 0.05	4.30	< 0.05	82	6.0	< 2.0	10.0	4.8	
	17-Jul	< 0.10	0.008	0.400	< 0.05	0.23	< 0.05	80	2.0	< 1.0	10.0	3.2	
	07-Aug	< 0.10	< 0.100	0.700	< 0.05	0.42	< 0.05	38	1.0	< 2.0	14.0	1.7	
	25-Sep	< 0.10	< 0.100	0.500	< 0.05	< 0.20	< 0.05	105	4.0	2.0	16.0	0.9	
	24-Oct	< 0.10	< 0.500	< 0.500	0.16	< 1.00	0.07	64	< 1.0	< 2.0	10.0	2.6	
	Mean		0.13	0.103	0.429	0.07	0.94	0.05	70	3.4	2.1	11.1	2.4
	Maximum		0.30	0.500	0.700	0.16	4.30	0.07	105	8.0	3.0	16.0	4.8
	Minimum		0.10	0.005	0.100	0.05	0.10	0.05	28	1.0	1.0	8.0	0.9
Std. Dev		0.08	0.180	0.250	0.04	1.51	0.01	29	2.7	0.7	2.8	1.3	
No. of D		1	1	5	1	4	1	7	6	1	7	7	
BZ-2S	24-Apr	0.20	< 0.005	< 0.100	< 0.05	0.10	< 0.05	< 10	14.0	< 3.0	6.0	0.6	
	24-May	0.10	< 0.005	0.400	< 0.05	< 0.10	< 0.05	72	< 1.0	< 3.0	6.0	0.2	
	14-Jun	< 0.10	< 0.005	< 0.100	< 0.05	5.40	< 0.05	58	9.0	< 2.0	8.0	2.1	
	17-Jul	0.10	< 0.005	0.300	< 0.05	0.45	< 0.05	84	3.0	< 1.0	10.0	0.2	
	07-Aug	< 0.10	< 0.100	0.300	1.20	0.74	< 0.05	28	6.0	< 2.0	10.0	0.3	
	25-Sep	0.10	< 0.100	0.200	0.07	< 0.20	< 0.05	66	4.0	1.0	10.0	0.8	
	24-Oct	< 0.10	< 0.500	< 0.500	0.07	< 1.00	< 0.05	56	< 1.0	< 2.0	10.0	0.1	
	Mean		0.11	0.103	0.271	0.22	1.14	0.05	53	5.4	2.0	8.6	0.6
	Maximum		0.20	0.500	0.500	1.20	5.40	0.05	84	14.0	3.0	10.0	2.1
	Minimum		0.10	0.005	0.100	0.05	0.10	0.05	10	1.0	1.0	6.0	0.1
Std. Dev		0.04	0.181	0.150	0.43	1.91	0.00	26	4.7	0.8	1.9	0.7	
No. of D		4	0	4	3	4	0	6	5	1	7	7	
BZ-3S	24-Apr	0.10	< 0.005	0.100	< 0.05	0.20	< 0.05	18	7.0	< 3.0	8.0	2.9	
	24-May	0.10	0.005	0.600	< 0.05	< 0.10	< 0.05	70	3.0	< 3.0	6.0	1.1	
	14-Jun	< 0.10	0.006	< 0.100	< 0.05	5.70	< 0.05	72	5.0	< 2.0	9.0	2.4	
	17-Jul	< 0.10	< 0.005	0.400	< 0.05	0.58	< 0.05	92	9.0	< 1.0	11.0	3.2	
	07-Aug	< 0.10	< 0.100	0.400	< 0.05	1.18	< 0.05	16	7.0	< 2.0	11.0	4.2	
	25-Sep	< 0.10	< 0.100	0.200	< 0.05	< 0.20	< 0.05	52	1.0	3.0	12.0	12.3	
	24-Oct	< 0.10	< 0.500	< 0.500	0.06	< 1.00	< 0.05	80	< 1.0	< 2.0	12.0	3.0	
	Mean		0.10	0.103	0.329	0.05	1.28	0.05	57	4.7	2.3	9.9	4.2
	Maximum		0.10	0.500	0.600	0.06	5.70	0.05	92	9.0	3.0	12.0	12.3
	Minimum		0.10	0.005	0.100	0.05	0.10	0.05	16	1.0	1.0	6.0	1.1
Std. Dev		0.00	0.181	0.198	0.00	1.99	0.00	30	3.1	0.8	2.3	3.7	
No. of D		2	2	5	1	4	0	7	6	1	7	7	

Table 3-2. (Continued)													
STATION	DATE	NH3	NO2	NO3	PO4	TKN	TP	TDS	TSS	BOD5	ALK	CHL A	
BZ-3M	24-Apr	< 0.10	< 0.005	0.100	< 0.05	0.40	< 0.05	< 10	3.0	< 3.0	8.0	2.7	
	24-May	< 0.10	< 0.005	0.600	< 0.05	< 0.10	< 0.05	76	7.0	< 3.0	6.0	1.0	
	14-Jun	< 0.10	0.005	< 0.100	< 0.05	3.40	< 0.05	74	5.0	< 2.0	9.0	8.5	
	17-Jul	< 0.10	0.005	0.500	< 0.05	0.53	< 0.05	104	5.0	< 1.0	12.0	1.2	
	07-Aug	< 0.10	< 0.100	0.700	< 0.05	0.53	< 0.05	34	< 1.0	< 2.0	10.0	1.4	
	25-Sep	< 0.10	< 0.100	0.500	0.06	< 0.20	< 0.05	61	4.0	1.0	12.0	0.8	
	24-Oct	< 0.10	< 0.500	< 0.500	0.08	< 1.00	< 0.05	84	< 1.0	< 2.0	16.0	2.1	
	Mean		0.10	0.103	0.429	0.06	0.88	0.05	63	3.7	2.0	10.4	2.5
	Maximum		0.10	0.500	0.700	0.08	3.40	0.05	104	7.0	3.0	16.0	8.5
	Minimum		0.10	0.005	0.100	0.05	0.10	0.05	10	1.0	1.0	6.0	0.8
Std. Dev		0.00	0.181	0.236	0.01	1.15	0.00	32	2.2	0.8	3.3	2.7	
No. of D		0	2	5	2	4	0	6	5	1	7	7	
BZ-3B	24-Apr	0.10	< 0.005	0.100	< 0.05	0.20	< 0.05	< 10	8.0	< 3.0	9.0	1.6	
	24-May	< 0.10	0.005	0.600	< 0.05	< 0.10	< 0.05	76	8.0	< 3.0	12.0	0.5	
	14-Jun	< 0.10	0.005	< 0.100	< 0.05	1.90	< 0.05	74	11.0	< 2.0	9.0	0.9	
	17-Jul	< 0.10	0.008	0.300	< 0.05	0.37	< 0.05	90	8.0	< 1.0	11.0	0.8	
	07-Aug	< 0.10	< 0.100	0.700	< 0.05	0.36	< 0.05	42	4.0	< 2.0	12.0	0.8	
	25-Sep	< 0.10	< 0.100	0.500	0.05	< 0.20	< 0.05	65	11.0	1.0	14.0	0.8	
	24-Oct	< 0.10	< 0.500	< 0.500	0.07	< 1.00	< 0.05	70	36.0	< 2.0	12.0	0.8	
	Mean		0.10	0.103	0.400	0.05	0.59	0.05	61	12.3	2.0	11.3	0.9
	Maximum		0.10	0.500	0.700	0.07	1.90	0.05	90	36.0	3.0	14.0	1.6
	Minimum		0.10	0.005	0.100	0.05	0.10	0.05	10	4.0	1.0	9.0	0.5
Std. Dev		0.00	0.180	0.238	0.01	0.65	0.00	27	10.7	0.8	1.8	0.3	
No. of D		10	3	5	2	4	0	6	7	1	7	7	
BZ-4S	24-Apr	0.10	< 0.005	< 0.100	< 0.05	0.40	< 0.05	10	5.0	< 3.0	5.0	1.9	
	24-May	< 0.10	< 0.005	0.300	< 0.05	0.10	< 0.05	54	4.0	< 3.0	10.0	11.7	
	14-Jun	< 0.10	< 0.005	< 0.100	< 0.05	5.90	< 0.05	42	4.0	< 2.0	7.0	1.1	
	17-Jul	< 0.10	0.005	0.400	< 0.05	0.43	< 0.05	80	1.0	< 2.0	8.0	0.8	
	07-Aug	< 0.10	< 0.100	0.200	< 0.05	0.38	< 0.05	40	9.0	6.0	10.0	1.3	
	25-Sep	< 0.10	< 0.100	0.500	< 0.05	< 0.20	< 0.05	52	7.0	5.0	18.0	0.7	
	24-Oct	< 0.10	< 0.500	< 0.500	0.07	< 1.00	< 0.05	72	< 1.0	< 2.0	10.0	0.9	
	Mean		0.10	0.103	0.300	0.05	1.20	0.05	50	4.4	3.3	9.7	2.6
	Maximum		0.10	0.500	0.500	0.07	5.90	0.05	80	9.0	6.0	18.0	11.7
	Minimum		0.10	0.005	0.100	0.05	0.10	0.05	10	1.0	2.0	5.0	0.7
Std. Dev		0.00	0.181	0.173	0.01	2.09	0.00	23	2.9	1.6	4.1	4.0	
No. of D		1	1	4	1	5	0	7	6	2	7	7	

Table 3-2. (Continued)													
STATION	DATE	NH3	NO2	NO3	PO4	TKN	TP	TDS	TSS	BOD5	ALK	CHL A	
BZ-5S	24-Apr	< 0.10	< 0.005	0.200	0.06	0.20	< 0.05	82	10.0	< 3.0	11.0	2.0	
	24-May	0.10	0.025	1.000	< 0.05	< 0.10	< 0.05	82	6.0	< 3.0	10.0	1.2	
	14-Jun	0.10	0.007	0.100	< 0.05	2.90	< 0.05	72	13.0	< 2.0	6.0	1.1	
	17-Jul	< 0.10	0.011	0.500	0.11	0.40	< 0.05	86	6.0	< 2.0	13.0	14.8	
	07-Aug	< 0.10	< 0.100	1.200	0.07	0.40	< 0.05	44	< 1.0	< 2.0	13.0	1.0	
	25-Sep	< 0.10	< 0.100	0.800	0.28	< 0.20	0.10	89	73.0	1.0	20.0	3.1	
	24-Oct	< 0.10	< 0.500	0.800	0.07	< 1.00	< 0.05	92	< 1.0	< 2.0	14.0	0.6	
	Mean		0.10	0.107	0.657	0.10	0.74	0.06	78	15.7	2.1	12.4	3.4
	Maximum		0.10	0.500	1.200	0.28	2.90	0.10	92	73.0	3.0	20.0	14.8
	Minimum		0.10	0.005	0.100	0.05	0.10	0.05	44	1.0	1.0	6.0	0.6
Std. Dev		0.00	0.178	0.408	0.08	1.00	0.02	16	25.6	0.7	4.3	5.1	
No. of D		2	3	7	5	4	1	7	5	1	7	7	
BZ-6S	24-Apr	< 0.10	< 0.005	0.100	< 0.05	0.30	< 0.05	86	1.0	< 3.0	5.0	1.9	
	24-May	< 0.10	0.006	0.600	< 0.05	< 0.10	< 0.05	90	6.0	< 3.0	12.0	2.0	
	14-Jun	< 0.10	0.007	< 0.100	< 0.05	9.20	< 0.05	36	10.0	< 2.0	9.0	2.5	
	17-Jul	< 0.10	0.006	0.400	< 0.05	0.43	< 0.05	92	1.0	< 2.0	10.0	5.4	
	07-Aug	< 0.10	< 0.100	0.400	< 0.05	0.53	< 0.05	28	4.0	< 2.0	11.0	3.6	
	25-Sep	< 0.10	< 0.100	0.200	< 0.05	< 0.20	< 0.05	53	2.0	< 1.0	10.0	8.8	
	24-Oct	< 0.10	< 0.500	< 0.500	0.07	< 1.00	< 0.05	66	< 1.0	< 2.0	12.0	3.9	
	Mean		0.10	0.103	0.329	0.05	1.68	0.05	64	3.6	2.1	9.9	4.0
	Maximum		0.10	0.500	0.600	0.07	9.20	0.05	92	10.0	3.0	12.0	8.8
	Minimum		0.10	0.005	0.100	0.05	0.10	0.05	28	1.0	1.0	5.0	1.9
Std. Dev		0.00	0.180	0.198	0.01	3.33	0.00	26	3.4	0.7	2.4	2.4	
No. of D		0	3	5	1	4	0	7	6	0	7	7	
BZ-6M	24-Apr	< 0.10	< 0.005	0.500	< 0.05	0.30	< 0.05	104	< 1.0	< 3.0	8.0	1.8	
	24-May	< 0.10	< 0.005	0.700	< 0.05	< 0.10	< 0.05	92	3.0	< 3.0	10.0	0.8	
	14-Jun	0.20	< 0.005	< 0.100	< 0.05	2.90	< 0.05	32	15.0	< 2.0	10.0	2.9	
	17-Jul	< 0.10	0.008	0.600	< 0.05	0.34	< 0.05	110	6.0	< 2.0	10.0	2.0	
	07-Aug	0.28	< 0.100	0.700	< 0.05	0.30	< 0.05	22	< 1.0	< 2.0	12.0	1.9	
	25-Sep	< 0.10	< 0.100	0.600	< 0.05	< 0.20	< 0.05	51	3.0	< 1.0	12.0	9.2	
	24-Oct	< 0.10	< 0.500	< 0.500	0.08	< 1.00	< 0.05	56	2.0	< 2.0	10.0	2.4	
	Mean		0.14	0.103	0.529	0.05	0.73	0.05	67	4.4	2.1	10.3	3.0
	Maximum		0.28	0.500	0.700	0.08	2.90	0.05	110	15.0	3.0	12.0	9.2
	Minimum		0.10	0.005	0.100	0.05	0.10	0.05	22	1.0	1.0	8.0	0.8
Std. Dev		0.07	0.180	0.206	0.01	1.00	0.00	35	5.0	0.7	1.4	2.8	
No. of D		2	1	5	1	4	0	7	5	0	7	7	

Table 3-2. (Continued)													
STATION	DATE	NH3	NO2	NO3	PO4	TKN	TP	TDS	TSS	BOD5	ALK	CHL A	
BZ-6B	24-Apr	< 0.10	< 0.005	0.200	< 0.05	0.30	< 0.05	80	3.0	< 3.0	10.0	1.3	
	24-May	< 0.10	0.005	0.600	< 0.05	< 0.10	< 0.05	80	2.0	< 3.0	10.0	0.6	
	14-Jun	0.10	< 0.005	< 0.100	< 0.05	4.10	< 0.05	44	20.0	< 2.0	9.0	1.1	
	17-Jul	< 0.10	0.010	0.300	< 0.05	0.11	< 0.05	38	4.0	< 2.0	11.0	1.2	
	07-Aug	0.12	< 0.100	0.700	0.06	0.28	< 0.05	22	1.0	< 2.0	12.0	1.0	
	25-Sep	< 0.10	< 0.100	0.400	2.43	< 0.20	1.44	62	10.0	5.0	12.0	1.4	
	24-Oct	< 0.10	< 0.500	< 0.500	0.07	< 1.00	< 0.05	78	< 1.0	< 2.0	12.0	0.5	
	Mean		0.10	0.104	0.400	0.39	0.87	0.25	58	5.9	2.7	10.9	1.0
	Maximum		0.12	0.500	0.700	2.43	4.10	1.44	80	20.0	5.0	12.0	1.4
	Minimum		0.10	0.005	0.100	0.05	0.10	0.05	22	1.0	2.0	9.0	0.5
Std. Dev		0.01	0.180	0.216	0.90	1.46	0.53	23	7.0	1.1	1.2	0.3	
No. of D		2	2	5	3	4	1	7	6	1	7	7	
BZ-7S	24-Apr	< 0.10	< 0.005	0.100	0.06	0.20	< 0.05	74	8.0	< 3.0	5.0	1.4	
	24-May	< 0.10	0.007	0.600	< 0.05	< 0.10	< 0.05	84	3.0	< 3.0	10.0	4.8	
	14-Jun	0.20	0.006	< 0.100	< 0.05	5.00	< 0.05	40	8.0	< 2.0	9.0	3.3	
	17-Jul	< 0.10	0.006	0.400	< 0.05	0.11	< 0.05	44	11.0	< 2.0	14.0	5.4	
	07-Aug	< 0.10	< 0.100	0.300	0.05	0.41	< 0.05	18	12.0	< 2.0	14.0	5.6	
	25-Sep	< 0.10	< 0.100	0.200	0.06	< 0.20	< 0.05	47	2.0	< 1.0	10.0	5.9	
	24-Oct	< 0.10	< 0.500	< 0.500	0.06	< 1.00	< 0.05	46	< 1.0	< 2.0	10.0	4.0	
	Mean		0.11	0.103	0.314	0.05	1.00	0.05	50	6.4	2.1	10.3	4.3
	Maximum		0.20	0.500	0.600	0.06	5.00	0.05	84	12.0	3.0	14.0	5.9
	Minimum		0.10	0.005	0.100	0.05	0.10	0.05	18	1.0	1.0	5.0	1.4
Std. Dev		0.04	0.180	0.195	0.01	1.79	0.00	22	4.4	0.7	3.1	1.6	
No. of D		1	3	5	4	4	0	7	6	0	7	7	
BZ-7M	24-Apr	0.20	< 0.005	0.100	< 0.05	0.30	< 0.05	92	3.0	< 3.0	8.0	3.4	
	24-May	< 0.10	0.007	0.700	< 0.05	< 0.10	< 0.05	88	3.0	< 3.0	12.0	3.1	
	14-Jun	< 0.10	0.006	< 0.100	0.06	4.10	< 0.05	34	6.0	< 2.0	10.0	6.0	
	17-Jul	< 0.10	0.011	0.500	< 0.05	0.34	< 0.05	34	7.0	< 2.0	12.0	1.1	
	07-Aug	< 0.10	< 0.100	0.500	0.08	0.47	< 0.05	32	11.0	< 2.0	16.0	3.5	
	25-Sep	< 0.10	< 0.100	0.200	0.05	< 0.20	< 0.05	46	2.0	6.0	12.0	5.9	
	24-Oct	< 0.10	< 0.500	< 0.500	0.09	< 1.00	< 0.05	58	1.0	< 2.0	12.0	3.9	
	Mean		0.11	0.104	0.371	0.06	0.93	0.05	55	4.7	2.9	11.7	3.8
	Maximum		0.20	0.500	0.700	0.09	4.10	0.05	92	11.0	6.0	16.0	6.0
	Minimum		0.10	0.005	0.100	0.05	0.10	0.05	32	1.0	2.0	8.0	1.1
Std. Dev		0.04	0.180	0.236	0.02	1.43	0.00	26	3.5	1.5	2.4	1.7	
No. of D		1	3	5	4	4	0	7	7	1	7	7	

Table 3-2. (Continued)												
STATION	DATE	NH3	NO2	NO3	PO4	TKN	TP	TDS	TSS	BOD5	ALK	CHL A
BZ-7B	24-Apr	< 0.10	< 0.005	0.100	0.06	0.20	< 0.05	84	8.0	< 3.0	10.0	1.2
	24-May	0.20	0.005	0.600	< 0.05	< 0.10	< 0.05	84	1.0	< 3.0	12.0	0.4
	14-Jun	< 0.10	0.005	< 0.100	0.07	4.60	< 0.05	34	11.0	< 2.0	10.0	1.5
	17-Jul	< 0.10	0.008	0.400	< 0.05	0.26	< 0.05	28	11.0	< 2.0	14.0	0.5
	07-Aug	0.11	< 0.100	0.600	0.09	0.36	< 0.05	46	5.0	< 2.0	14.0	1.0
	25-Sep	< 0.10	< 0.100	0.400	0.05	< 0.20	< 0.05	63	6.0	2.0	14.0	1.9
	24-Oct	< 0.10	< 0.500	< 0.500	0.08	< 1.00	< 0.05	68	6.0	< 2.0	12.0	4.2
Mean		0.12	0.103	0.386	0.06	0.96	0.05	58	6.9	2.3	12.3	1.5
Maximum		0.20	0.500	0.600	0.09	4.60	0.05	84	11.0	3.0	14.0	4.2
Minimum		0.10	0.005	0.100	0.05	0.10	0.05	28	1.0	2.0	10.0	0.4
Std. Dev		0.04	0.180	0.212	0.02	1.63	0.00	23	3.5	0.5	1.8	1.3
No. of D		2	3	5	5	4	0	7	7	1	7	7

Ammonia




Figure 3-11. Ammonia measured in surface, middle, and bottom waters of Beltzville Reservoir in 2001.

reservoirs usually results in those that are affected by eutrophication. However, high concentrations of ammonia were not observed on later monitoring dates in September and October when hypoxia was present in the lower water column.

Concentrations of ammonia measured at Beltzville Reservoir were in compliance with the PADEP water quality standards during 2001. The state water quality standard for ammonia is dependent on temperature and pH (Table 3-3). Throughout the monitoring period, the most conservative criterion for ammonia resulting from the highest measured pH (8.6) and temperature (30 °C) was 0.6-mg/L.

pH	10°C	15°C	20°C	25°C	30°C
6.50	25.5	17.4	12.0	8.4	5.9
6.75	23.6	16.0	11.1	7.7	5.5
7.00	20.6	14.0	9.7	6.8	4.8
7.25	16.7	11.4	7.8	5.5	3.9
7.50	12.4	8.5	5.9	4.1	2.9
7.75	8.5	5.8	4.0	2.8	2.0
8.00	5.5	5.8	4.0	2.8	2.0
8.25	3.4	2.3	1.6	1.2	0.9
8.50	2.0	1.4	1.0	0.7	0.6
8.75	1.2	0.9	0.6	0.5	0.4
9.00	0.8	0.5	0.4	0.3	0.3
9.25	0.36	0.24	0.17	0.12	0.08
9.50	0.20	0.13	0.10	0.07	0.05

Seasonal trends for ammonia in surface water were determined for individual stations using the non-parametric Mann-Kendall statistic. The statistical analysis was conducted for stations with historical data (21 years or more), individually for spring (April through June) and summer (July through September) seasons. Data from seven stations were analyzed representing downstream (BZ-1), Pine Run (BZ-2), the reservoir-body (BZ-3, -6, and -7), Wild Creek (BZ-4), and upstream on Pohopoco Creek (BZ-5). Ammonia concentrations appear to be decreasing throughout the Beltzville Reservoir watershed (Table 3-4). From the analysis, stations BZ-1, -2, -3, -4, and -5 showed significant and decreasing trends. Rates of decrease were greater during the summer season and ranged from 0.001 to 0.004 mg/L/year. Stations BZ-1 and -5 also had significant decreasing spring season rates of 0.001 mg/L/year.

Table 3-4. Seasonal trends of ammonia concentration at individual stations of Beltzville Reservoir calculated with the Mann-Kendall Statistic. Shaded values are significant (at least $P < 0.05$).

Station	# of Years spring/summer	Spring		Summer	
		P Level	Rate (mg/)	P Level	Rate (mg/L)
Surface Water					
BZ-1	24/23	<0.05	-0.001	<0.001	-0.004
BZ-2	22	NS	-0.001	<0.05	-0.001
BZ-3	26/25	NS	-0.001	<0.01	-0.003
BZ-4	25/24	NS	-0.001	<0.01	-0.002
BZ-5	25	<0.001	-0.001	<0.001	-0.003
BZ-6	6	NS	0.009	NS	0.005
BZ-7	6	NS	0.005	NS	0.008

3.2.2 Nitrite and Nitrate

Nitrite concentrations in the water column of Beltzville Reservoir were very low during 2001 (Fig. 3-12). Concentrations measured at all stations and depths were less than the method detection limit of 0.5-mg/L throughout the monitoring period.

Nitrate was distributed uniformly in the water column of Beltzville Reservoir during 2001 (Fig. 3-13). At most stations and depths, concentrations ranged from less than the method detection limit (0.1-mg/L) to 0.8-mg/L. Concentrations in surface water from the reservoir-body stations (BZ-3, -6, and -7) were similar to those from middle and bottom depths. Overall, concentrations appeared to average 0.4-mg/L throughout the monitoring period. Concentrations of nitrate upstream of the reservoir on Pohopoco Creek (BZ-5) were usually highest and over the monitoring period ranged from 0.1 to 1.2-mg/L. Nitrate measured downstream of the reservoir (BZ-1) was consistent over the monitoring period at about 0.4-mg/L. This average was consistent with all other stations.

Beltzville Reservoir was in compliance with the PADEP water quality standard for nitrite and nitrate during 2001. The standard is a summed concentration of nitrite and nitrate of less than 10 mg/L. Throughout the monitoring period, the summed concentrations for all stations averaged 0.5-mg/L with maximum values of 1.3-mg/L in August and October at BZ-5

3.2.3 Total Inorganic Nitrogen

Concentrations of total inorganic nitrogen have not changed appreciably over the past 26 years in surface waters of Beltzville Reservoir or downstream of the reservoir in either spring or summer seasons (Figs. 3-14 and 3-15).




Figure 3-12. Nitrite measured in surface, middle, and bottom waters of Beltzville Reservoir in 2001.





Figure 3-13. Nitrate measured in surface, middle, and bottom waters of Beltville Reservoir in 2001.


Total Nitrogen *Spring*



3-25

Figure 3-14. Seasonal trends of total nitrogen (ammonia + nitrite + nitrate) in the spring at Beltzville Reservoir.

Total Nitrogen Summer



3-26

Figure 3-15. Seasonal trends of total nitrogen (ammonia + nitrite + nitrate) in the summer at Beltzville Reservoir.

Seasonal trends (spring and summer) for total nitrogen in surface water were also determined for individual stations using the non-parametric Mann-Kendall statistic. Total inorganic nitrogen, calculated as the summed concentration of ammonia, nitrite, and nitrate, was analyzed separately for reservoir and upstream stations (BZ-2, -3, -4, -5, -6, and -7) and downstream of the reservoir (station BZ-1). Regression analyses conducted on the average concentrations indicated significant decreasing trends for stations BZ-1, -3, and -4 during the summer period. Throughout the 26-year time series, total nitrogen concentrations generally averaged less than 0.8-mg/L; however, in 2001, concentrations were slightly higher at about 1.0-mg/L.

Table 3-5. Seasonal trends of total nitrogen at individual stations of Beltzville Reservoir calculated with the Mann-Kendall Statistic. Shaded values are significant at P – 0.05.

Station	# of Years Spring/Summer	Spring		Summer	
		P Level	Rate (mg/L)	P Level	Rate
Surface Water					
BZ-1	24/23	NS	-0.004	< 0.01	-0.01
BZ-2	22	NS	-0.002	NS	0.002
BZ-3	26/25	NS	-0.002	< 0.05	-0.08
BZ-4	25/24	NS	-0.004	< 0.05	-0.01
BZ-5	25	NS	-0.007	NS	0.03
BZ-6	6	NS	-0.03	NS	0.02
BZ-7	6	NS	-0.05	NS	0.008

3.2.4 Total Kjeldahl Nitrogen

For the most part, total Kjeldahl nitrogen (TKN) was uniformly low in the water column of Beltzville Reservoir during 2001 (Fig. 3-16). Throughout the monitoring period, concentrations measured at all depths generally ranged less than 1.2-mg/L. In June; however, concentrations averaged 2.3-mg/L and ranged as high as 4.6-mg/L. Concentrations measured at upstream stations BM-2, -4, and -5 were 2.7, 2.95, and 1.45-mg/L, respectively. Concentrations were also elevated within the reservoir body at stations BZ-3, -6, and -7 and below the reservoir at station BZ-1. Concentrations at these stations measured 2.85, 4.6, 2.5, and 2.15-mg/L, respectively.

Total Kjeldahl Nitrogen




Figure 3-16. Total Kjeldahl nitrogen measured in surface, middle, and bottom waters of Beltzville Reservoir in 2001.

3.2.5 Dissolved Phosphate

Dissolved phosphate was not a significant nutrient parameter at Beltzville Reservoir in 2001 (Fig. 3-17). Concentrations of dissolved phosphate were generally at or below the detection limit of 0.05-mg/L. Two isolated instances of high values were recorded during August at BZ-2 in the surface water and in September at BZ-6 in the bottom water. These concentrations were 1.2 and 2.4-mg/L, respectively. In freshwater environments, dissolved phosphate is usually a limiting nutrient and is readily taken up by freshwater plants and algae.

3.2.6 Total Phosphorus

Total phosphorus in the water column of Beltzville Reservoir was consistently measured at low concentrations during 2001 (Fig. 3-18). EPA guidance for nutrient criteria in lakes and reservoirs suggests a minimum concentration for total phosphorus of 0.01-mg/L (EPA 2000). Lakes and reservoirs exceeding this concentration are more likely to experience algal bloom problems during the growing season. Overall, only three measures for total phosphorus with results greater than the detection limit from reservoir monitoring were greater than the EPA guideline. High measures of total phosphorus were recorded in the surface waters at station BZ-5 (0.1-mg/L) in September and BZ-1 (0.7-mg/L) in October. Additionally, a high concentration of total phosphorus was present in the bottom water at BZ-6 (1.4-mg/L) in September. All remaining results were less than the detection limit of 0.05-mg/L, which is a concentration 5 times greater than the guideline.

Total phosphorus concentrations measured during 2001 and historical data from the past 26 years were analyzed for seasonal trends. Regression analyses for spring and summer periods were conducted separately for stations of the reservoir and downstream. No trends were determined for either season at reservoir and downstream stations (Figs. 3-19 and 3-20). None of the regression lines were significant ($P > 0.05$).

Trend analyses for total phosphorus were also conducted for individual monitoring stations of Beltzville Reservoir using the non-parametric Mann-Kendall statistic. As before, the trends were calculated separately for spring and summer seasons. Stations analyzed for the seasonal trends included BZ-1, -2, -3, -4, -5, -6, and -7. From these stations, historical data on total phosphorus spanned a maximum of 21 years. Only one significant trend resulted from the analysis. A significant decreasing trend in summer was determined for station BZ-4 on the Wild Creek tributary (Table 3-6). The estimated rate of decrease for this trend was 0.004-mg/L/year.

Dissolved Phosphate




Figure 3-17. Dissolved phosphorus measured in surface, middle, and bottom waters of Beltzville Reservoir in 2001.

Total Phosphorus





Figure 3-18. Total phosphorus measured in surface, middle, and bottom waters of Beltzville Reservoir in 2001.

Total Phosphorus Spring



3-32

Figure 3-19. Seasonal trends of total phosphorus in spring at Beltzville Reservoir.

Total Phosphorus *Summer*




Figure 3-20. Seasonal trends of total phosphorus in summer at Beltzville Reservoir.

Table 3-6. Seasonal trends of total phosphorus concentration at individual stations of Beltzville Reservoir calculated with the Mann-Kendall Statistic. Shaded values are significant (at least $P < 0.05$).

Station	# of Years spring/summer	Spring		Summer	
		P Level	Rate (mg/L)	P Level	Rate (mg/L)
Surface Water					
BZ-1	19	NS	0.0	NS	0.0003
BZ-2	21	NS	0.09	NS	-0.0004
BZ-3	22/21	NS	-0.13	NS	-0.0003
BZ-4	21/20	NS	0.28	< 0.05	-0.004
BZ-5	21	NS	-0.09	NS	-0.00001
BZ-6	5	NS	5.75	NS	0.007
BZ-7	5	NS	6.44	NS	0.006

3.2.7 Total Dissolved Solids

Concentrations of total dissolved solids (TDS) in the water column of Beltzville Reservoir were variable during 2001 (Fig. 3-21). The variability among measures was different on each monitoring date, and patterns of consistency for individual stations were not easily discernible. Concentrations among all stations and depths ranged from 10 to 104-mg/L. Overall, measures of TDS at all stations and depths averaged 60-mg/L.

TDS concentrations measured at Beltzville Reservoir in 2001 were in compliance with PADEP water quality standards. The state water quality standard for TDS is a maximum concentration of 500 mg/L. Throughout the monitoring period, all measures of TDS were at least 5 times less than the standard (Fig. 3-21).

Total dissolved solids concentrations have exhibited no discernible pattern over the past 26 years of water quality monitoring at Beltzville Reservoir. Regression analyses conducted separately by season for upstream and downstream data did not result in any significant trends (Figs. 3-22 and 3-23). Average TDS concentrations calculated this year were slightly higher than recent years.

Seasonal trends (spring and summer) for total dissolved solids in surface water were also determined for individual stations using the non-parametric Mann-Kendall statistic. Overall, seven stations were tested representing downstream (BZ-1), Pine Run (BZ-2), mid-reservoir (BZ-3, -6, and -7), Wild Creek (BZ-4), and upstream on Pohopoco Creek (BZ-5). Concentrations of total dissolved solids appear to have decreased in many of the surface waters of Beltzville Reservoir over the past two decades (Table 3-7). Significant




Figure 3-21. Total dissolved solids in surface, middle, and bottom waters of Beltzville Reservoir in 2001. (The PADEP water quality standard for TDS is a maximum concentration of 500 mg/l.)

Total Dissolved Solids *Spring*





Figure 3-22. Seasonal trends of total dissolved solids in spring at Beltzville Reservoir.

Total Dissolved Solids *Summer*



3-37

Figure 3-23. Seasonal trends of total dissolved solids in summer at Beltzville Reservoir.

decreasing trends were observed in the mid-reservoir (station BZ-3) during both seasons as well as for Pine Creek and Wild Creek tributaries (stations BZ-2 and BZ-4) in the summer. The estimated rates of decrease among all trends were similar at about 0.8-mg/L/year. A significant increasing trend was seen at station BZ-6 during the spring period. The rate of increase was 11.5-mg/L/yr.

Table 3-7. Seasonal trends of total dissolved solids concentration at individual stations of Beltzville Reservoir calculated with the Mann-Kendall Statistic. Shaded values are significant (at least $P < 0.05$).

Station	# of Years spring/summer	Spring		Summer	
		P Level	Rate (mg/L)	P Level	Rate (mg/L)
Surface Water					
BZ-1	23	NS	-0.047	NS	-0.527
BZ-2	22	NS	-0.583	< 0.05	-1.02
BZ-3	25	< 0.05	-0.579	< 0.05	-0.742
BZ-4	24	NS	-0.342	< 0.05	-0.922
BZ-5	25	NS	0.105	NS	-0.487
BZ-6	6	< 0.05	11.5	NS	5
BZ-7	6	NS	5.5	NS	-2.33

3.2.8 Total Suspended Solids

Total suspended solids (TSS) concentrations in the water column of Beltzville Reservoir were generally low during 2001 (Fig. 3-24). In general, concentrations measured at most stations and depths ranged less than 20 mg/L. Higher concentrations were measured in September at the surface of station BZ-5 and in the bottom waters of station BZ-3 in October. The concentration of TSS at BZ-5 was 73-mg/L and 36-mg/L at BZ-3.

3.2.9 Biochemical Oxygen Demand

Biochemical oxygen demand (BOD) is a measure of the oxygen-depleting burden imposed by organic material present in water. BOD concentrations in the water column of Beltzville Reservoir were relatively low in 2001 (Fig. 3-25). Throughout the monitoring period, concentrations measured at 91% of all stations and depths were typically less than the method detection limit (1 to 3-mg/L). BOD was detected at all stations and throughout the water column in September and only once in August on Wild Creek (BZ-4). Concentrations at all stations and depths in the water column averaged 2.3-mg/L throughout the monitoring period.

Total Suspended Solids




Figure 3-24. Total suspended solids in surface, middle, and bottom waters of Beltzville Reservoir in 2001.

Biochemical Oxygen Demand




Figure 3-25. Biochemical oxygen demand in surface, middle, and bottom waters of Beltzville Reservoir in 2001.

Based on trend analyses using 2001 and historical data, BOD does not appear to have changed significantly in the reservoir or downstream (Figs. 3-26 and 3-27). Although the regression lines for both spring and summer seasons suggested a slight reduction in BOD concentrations, none of the trends were significant ($P > 0.05$).

A trend analysis for BOD was also conducted on individual monitoring stations of Beltzville Reservoir using the non-parametric Mann-Kendall statistic (Table 3-8). Based on this analysis, a significant decreasing summer trend was determined for station BZ-1, downstream of the reservoir. The estimated rate of decrease ranged was 0.03-mg/L/year.

Table 3-8. Seasonal trends of total BOD concentration at individual stations of Beltzville Reservoir calculated with the Mann-Kendall Statistic. Shaded values are significant (at least $P < 0.05$).					
Station	# of Years spring/summer	Spring		Summer	
		P Level	Rate (mg/L)	P Level	Rate (mg/L)
Surface Water					
BZ-1	19	NS	-0.003	<0.01	-0.03
BZ-2	21	NS	0.0	NS	-0.004
BZ-3	21	NS	-0.02	NS	-0.009
BZ-4	20	NS	0.0	NS	0.0
BZ-5	21	NS	0.0	NS	-0.04
BZ-6	6	NS	0.13	NS	0.0
BZ-7	6	NS	0.13	NS	0.0

3.2.10 Alkalinity

Alkalinity is a measure of the acid-neutralizing capacity of water. Alkalinity in the water column of Beltzville Reservoir was relatively low during 2001 (Fig. 3-28). Throughout the monitoring period, concentrations measured at all stations and depths ranged between 5 and 20-mg/L.

Concentrations of alkalinity measured at Beltzville Reservoir were in compliance with PADEP water quality standards during 2001. The state water quality standard for alkalinity is a minimum concentration of 20 mg/L CaCO_3 , except where natural conditions are less. The natural alkalinity of water is largely dependent on the underlying geology and soils within the surrounding watershed. The low alkalinity measured at Beltzville Reservoir probably results from the regional geology, which is primarily sandstone and shale (Van Diver 1990).

5-day Biochemical Oxygen Demand *Spring*

3-42




Figure 3-26. Seasonal trends of five-day biochemical oxygen demand in spring at Beltzville Reservoir.

5-day Biochemical Oxygen Demand *Summer*

3-43




Figure 3-27. Seasonal trends of five-day biochemical oxygen demand in summer at Beltzville Reservoir.

3.2.11 Total Organic and Inorganic Carbon

Total inorganic carbon and total organic carbon in the water column of Beltzville Reservoir were not present in measurable concentrations during 2001 (Fig. 3-29 and Fig. 3-30). All concentrations measured were less than the method detection limit of 5-mg/L.

3.2.12 Chlorophyll *a*

Chlorophyll *a* is a measure of algal biomass. For the most part, chlorophyll *a* concentrations in the water column of Beltzville Reservoir were relatively low during 2001 (Fig. 3-31). Concentrations measured at all stations and depths ranged between 0.1 and 14.8-mg/m³. Concentrations were highly variable throughout the reservoir and its tributaries and averaged 2.6- mg/m³.

3.2.13 BTEX

Concentrations of benzene, toluene, ethylbenzene, o-xylene and m,p-xylenes (BTEX) in surface waters of Beltzville Reservoir were consistently low during 2001 (Table 3-9). Benzene and Ethylbenzene concentrations were always less than method detection limits of 1.0 µg/L. Toluene was detected in April, June, and August with values ranging from 1 to 4 µg/L. o-Xylene concentrations were only detected one time in August with a low value of 1 µg/L. m,p-Xylene was detected in August at all three stations (BZ-3, -6, and -7) with values of 2 and 4 µg/L.

Beltzville Reservoir was in compliance with PADEP water quality standard for BTEX parameters during 2001. All of the concentrations measured were more than an order of magnitude less than PADEP water quality criteria for toxic substances. PADEP fish and aquatic criteria continuous concentrations (CCCs) for the 4 parameters are as follows: benzene, 128µg/L ; ethylbenzene, 580 µg/L; toluene, 330 µg/L; and xylene, 211 µg/L (Pa. Code, Title 25, Chapter 16).




Figure 3-28. Alkalinity in surface, middle, and bottom waters of Beltzville Reservoir in 2001. (The PADEP water quality standard for alkalinity is a minimum concentration of 20 mg/L, except where natural conditions are less.)




Figure 3-29. Total inorganic carbon in surface, middle, and bottom waters of Beltzville Reservoir in 2001.

Total Organic Carbon




Figure 3-30. Total organic carbon in surface, middle, and bottom waters of Beltzville Reservoir in 2001.




Figure 3-31. Total Chlorophyll a in surface, middle, and bottom waters of Beltzville Reservoir in 2001.

Station	Date	Benzene	Ethylbenzene	Toluene	o-Xylene	m,p-Xylene
BZ-3S	24-Apr	< 1	< 1	1	< 1	< 1
	24-May					
	14-June	< 1	< 1	2	< 1	1
	17-July	< 1	< 1	< 1	< 1	< 1
	7-Aug	< 1	< 1	3	1	4
	25-Sept	< 1	< 1	< 1	< 1	< 1
	24-Oct	< 1	< 1	< 1	< 1	< 1
BZ-6S	24-Apr	< 1	< 1	< 1	< 1	< 1
	24-May					
	14-June	< 1	< 1	4	< 1	< 1
	17-July	< 1	< 1	< 1	< 1	< 1
	7-Aug	< 1	< 1	2	< 1	2
	25-Sept	< 1	< 1	< 1	< 1	< 1
	24-Oct	< 1	< 1	< 1	< 1	< 1
BZ-7S	24-Apr	< 1	< 1	1	< 1	< 1
	24-May					
	14-June	< 1	< 1	< 1	< 1	< 1
	17-July	< 1	< 1	< 1	< 1	< 1
	7-Aug	< 1	< 1	2	< 1	2
	25-Sept	< 1	< 1	< 1	< 1	< 1
	24-Oct	< 1	< 1	< 1	< 1	< 1

* BTEX was not analyzed during May monitoring because of laboratory sample processing error

3.3 TROPHIC STATE DETERMINATION

Carlson's (1977) trophic state index (TSI) is a method of quantitatively expressing the magnitude of eutrophication for a lake. The trophic state analysis calculates separate indices for eutrophication based on measures of total phosphorus, chlorophyll *a*, and secchi disk. Index values for each parameter range on the same scale from 0 (least enriched) to 100 (most enriched). The resulting indices can also be compared to qualitative threshold values that correspond to levels of eutrophication: mesotrophic (TSI < 40), mesoeutrophic (TSIs from 50 to 60), and eutrophic (TSI > 60).

TSIs calculated for measures of secchi disk depth classified Beltzville Reservoir as both oligotrophic and mesotrophic during 2001 (Fig. 3-32). The TSI values ranged from 28 to 36 when it was oligotrophic and from 40 to 47 during the mesotrophic period.

Trophic State




Figure 3-32. Trophic state indices calculated from secchi disk depth and concentrations of total phosphorus and chlorophyll a at Beltzville Reservoir in 2001

The TSI classification based on measures of total phosphorus was not feasible because of the high method detection limit for the parameter (Fig. 3-32). All concentrations measured at Beltzville Reservoir were less than the method detection limit of 0.05-mg/L. However, at this concentration, the TSI result of 60 is already bordering on eutrophic.

TSIs calculated for measures of chlorophyll *a* classified Beltzville Reservoir as mesotrophic (Fig. 3-32). Most of the TSI values ranged from 40 to 45. There were two monitoring dates that were not in that range. On 24 April, the TSI value was 37.67, which is in the oligotrophic level and on 25 September the TSI value was 52.13, which is in the mesoeutrophic level.

Carlson (1977) warned against averaging TSI values estimated for different parameters, and instead suggested giving priority to chlorophyll *a* in the summer and to phosphorus in the spring, fall, and winter. With this in mind, our estimation of the trophic state of the reservoir based on TSI's was mesotrophic during 2001.

The EPA (1983) also provides criteria for defining the trophic conditions of lakes of the north-temperate zone based on concentrations of total phosphorus, chlorophyll *a*, and secchi depth (Table 3-10). As related above, total phosphorus was not measured with enough precision to permit a substantive interpretation of trophic condition. The concentration of the method detection limit used for total phosphorus was eutrophic. Concentrations of chlorophyll *a* and secchi disk depth ranged from oligotrophic (April to June and October) to mesotrophic (July to September). Taking into account the general agreement between the EPA classifications with that of the TSIs, the trophic condition of Beltzville Reservoir was borderline oligotrophic-mesotrophic.

Water Quality Variable	Oligo-trophic	Meso-trophic	Eutrophic	Apr	May	Jun	Jul	Aug	Sep	Oct
Total phos. ($\mu\text{g/l}$)	< 10	10-20	> 20	50	50	50	50	50	50	50
Chlorophyll (mg/m^3)	< 4	4-10	> 10	2.06	2.65	2.74	4.67	4.45	9.00	3.60
Secchi depth (m)	> 3.7	2-4	< 2	3.83	4.99	4.95	8.86	2.45	2.82	3.40

3.4 RESERVOIR BACTERIA MONITORING

Three forms of coliform bacteria contamination were monitored at Beltzville Reservoir during 2001 including total coliform, fecal coliform, and fecal streptococcus (Table 3-11). Total coliform includes *Escherichia coli* (*E. coli*) and related bacteria that are associated with fecal discharges. Fecal coliform bacteria are a subgroup of the total coliform and are normally associated with waste derived from human and other warm-blooded animals. Fecal streptococcus refers to a group of bacteria that are common to warm-blooded animals other than humans. Because of the relative differences between fecal coliform and fecal streptococcus, their ratio can be used as a qualitative indicator of the source of fecal contamination.

Total coliform contamination of Beltzville Reservoir was generally low during 2001 monitoring period (Fig. 3-33). No station appeared to have consistently higher counts over the monitoring period. However, all stations during the autumn season (September and October), had higher total coliform counts that ranged from 60 to 560-clns/100ml. The highest counts of 560-clns/100-ml were found at station BZ-1, on 25 September.

Fecal coliform counts were generally lower counts than total coliform except for station BZ-4 (Fig. 3-34). On 24 October monitoring, BZ-4 had the highest count of 850-clns/100-mls. Counts within the reservoir (stations BZ-3, BZ-6, and BZ-7) were low throughout the monitoring period.

Fecal streptococcus contamination at Beltzville Reservoir had low counts below 200-clns/100-mls except for one station BZ-2 (Fig. 3-35). On 7 August station BZ-2 had the highest count of 330-clns/100-mls observed during monitoring period. Counts within the reservoir (stations BZ-3, BZ-6, and BZ-7) and downstream (station BZ-1) were low generally ranging less than 120-clns/100-ml.

With respect to PADEP water quality standards, coliform bacteria contamination was low at Beltzville Reservoir during 2001 (Table 3-12; Fig. 3-36). The PADEP standard for bacteria during the swimming season (from 1 May to 30 September) is a geometric mean not greater than 200 colonies/100-ml calculated for fecal coliform samples collected on consecutive days. Given that our regular monitoring was completed on one day, we calculated the geometric mean based on all the surface water samples collected on that day. Throughout the monitoring period, the geometric mean, calculated for each monitoring day, was less than the PADEP criteria.

Flow data from a USGS gauging station within the Beltzville Reservoir watershed (Kresgeville) were analyzed to qualitatively correlate precipitation patterns with coliform bacteria contamination (Fig. 2-2 through 2-6). Overall, coliform contamination did not appear to be strongly correlated with precipitation and streamflow. The only time that precipitation may have effected coliform bacteria contamination is during the monitoring on




Figure 3-33. Total coliform counts in surface waters of Beltzville Reservoir in 2001

3-54




Figure 3-34. Fecal coliform counts in surface waters of Beltzville Reservoir in 2001. (PADEP water quality standard for fecal coliform is less than 200 colonies/100 mls.)

Fecal Streptococcus




Figure 3-35. Fecal streptococcus counts in surface waters of Beltzville Reservoir in 2001

Fecal Coliform

3-56




Figure 3-36. Maximum, average, and geometric mean of fecal coliform counts (colonies/100-ml) for all stations monitored at Beltzville Reservoir in 2001

25 September. This rain event exceeded four inches of rain and the total coliform counts are high ranging from 60 to 560-clns/100-mls.

Table 3-11. Bacteria counts (colonies/100 ml) at Beltzville Reservoir during 2001.

STATION	DATE	Total Coliform (TC)	Fecal Coliform (FC)	Fecal Strep (FS)	FC/FS
BZ-1S	24-Apr	60	< 10	1	NC
	24-May	20	< 10	< 10	NC
	14-Jun	70	< 10	10	NC
	17-Jul	50	30	10	3.00
	07-Aug	34	20	28	0.71
	25-Sep	560	220	< 10	NC
	24-Oct	200	200	< 10	NC
BZ-2S	24-Apr	180	< 10	10	NC
	24-May	30	10	< 10	NC
	14-Jun	140	180	40	4.50
	17-Jul	50	< 10	70	NC
	07-Aug	116	108	330	0.33
	25-Sep	220	100	170	0.59
	24-Oct	120	40	40	1.00
BZ-3S	24-Apr	10	10	< 1	NC
	24-May	< 10	< 10	< 10	NC
	14-Jun	< 10	< 10	< 10	NC
	17-Jul	20	< 10	< 10	NC
	07-Aug	6	< 2	4	NC
	25-Sep	360	120	40	3.00
	24-Oct	100	20	< 10	NC
BZ-4S	24-Apr	10	10	< 1	NC
	24-May	10	130	10	13.00
	14-Jun	50	50	30	1.67
	17-Jul	20	60	30	2.00
	07-Aug	24	74	54	1.37
	25-Sep	100	100	< 10	NC
	24-Oct	280	850	30	28.33
BZ-5S	24-Apr	320	< 10	1	NC
	24-May	240	200	140	1.43
	14-Jun	150	140	80	1.75
	17-Jul	10	< 10	10	NC
	07-Aug	290	226	136	1.66
	25-Sep	150	140	40	3.50
	24-Oct	300	50	60	0.83
	24-Oct	220	40	120	0.33

Table 3-11. (Continued)

STATION	DATE	Total Coliform (TC)	Fecal Coliform (FC)	Fecal Strep (FS)	FC/FS
BZ-6S	24-Apr	< 10	100	< 1	NC
	24-May	10	< 10	< 10	NC
	14-Jun	< 10	< 10	< 10	NC
	17-Jul	50	50	< 10	NC
	07-Aug	6	2	4	0.50
	25-Sep	200	20	< 10	NC
BZ-7S	24-Apr	50	< 10	< 1	NC
	24-May	< 10	< 10	< 10	NC
	14-Jun	< 10	10	< 10	NC
	17-Jul	40	< 10	< 10	NC
	07-Aug	6	2	6	0.33
	25-Sep	60	20	< 10	NC
	24-Oct	180	20	100	0.20

Table 3-12. Summary statistics of fecal coliform counts (colonies/100-ml) among all stations of Beltzville Reservoir during 2001. (PADEP water quality standard for fecal coliforms is a geometric mean not greater than 200 colonies/100-ml.)

	Geometric Mean	Arithmetic Mean	Maximum Count
24-Apr	13.9	32.5	100.0
24-May	22.1	48.8	200.0
14-Jun	27.7	52.5	180.0
17-Jul	19.0	28.8	60.0
07-Aug	16.2	54.5	226.0
25-Sep	76.1	92.5	220.0
24-Oct	66.0	157.5	850.0

An analysis for trends was conducted on fecal coliform data collected during 2001 and historical data collected over the past 21 years. Average counts at reservoir and downstream stations were analyzed separately, for trends occurring in spring and summer sessions. None of the regression lines were significant for either location in either season (Figs. 3-37 and 3-38). However, the plotted averages for both reservoir and downstream sites over the past 21-year time frame reflect consistently low fecal coliform contamination.

Seasonal trend analyses for total and fecal coliform contamination in surface water were also calculated for individual stations using the Mann-Kendall Statistic (Tables 3-13 and 3-14). The tests were conducted for spring (April through June) and summer (July through September) seasons for seven stations with extensive historical data. No trends were observed for total coliforms (Table 3-13). For fecal coliforms there were two trends that were




Figure 3-37. Seasonal trends of fecal coliform in spring at Beltzville Reservoir

Fecal Coliform *Summer*




Figure 3-38. Seasonal trends of fecal coliform in summer at Beltzville Reservoir

determined to be significant; station BZ-5 and BZ-6 for fecal coliform contaminants in the spring (Table 3-13). The rate of increase estimated for the trend was 8.58 and 3.0 colonies/100-ml/year.

Table 3-13. Seasonal trends of fecal coliforms/100-ml at individual stations of Beltzville Reservoir calculated with the Mann-Kendall Statistic.					
Station	# of Years spring/summer	Spring		Summer	
		P Level	Rate (mg/L)	P Level	Rate
Surface Water					
BZ-1	17/16	NS	-0.06	NS	8.06
BZ-2	21	NS	-4.75	NS	6.11
BZ-3	21/20	NS	-2.78	NS	-1.14
BZ-4	21/20	NS	-6.75	NS	-3.54
BZ-5	21	NS	-18.61	NS	3.64
BZ-6	6	NS	0.5	NS	3.67
BZ-7	6	NS	-2	NS	-72.33

Table 3-14. Seasonal trends of fecal coliforms/100-ml at individual stations of Beltzville Reservoir calculated with the Mann-Kendall Statistic.					
Station	# of Years spring/summer	Spring		Summer	
		P Level	Rate (mg/L)	P Level	Rate
Surface Water					
BZ-1	17	NS	0.12	NS	1.07
BZ-2	21	NS	0.82	NS	3.35
BZ-3	21	NS	-0.13	NS	-1.07
BZ-4	21	NS	-0.42	NS	0.82
BZ-5	21	< 0.05	8.58	NS	6.37
BZ-6	6	< 0.05	3.0	NS	3.25
BZ-7	6	NS	0.75	NS	-104

The ratio of fecal coliform to fecal streptococcus counts can be used to qualitatively identify sources of bacteria contamination (McComas 1993). The ratio is characteristic for several animal species and certain waste disposal practices; for human waste, the ratio is 4 to 1. Use of the ratio was limited to the extent that counts at most stations were at or less than method detection limits for one or both coliform parameters. Out of a total of 49 perspective ratios, only 21 were calculated. Of these, ratios ranged from 0.2 to 28.33, and only 10 fell within the range one would expect for human waste sources (Table 3-12).

3.5 SEDIMENT PRIORITY POLLUTANT MONITORING

Sediment samples were collected at station BZ-6 and analyzed for priority pollutant contaminants, Group 2 – semivolatile organic compounds and metals. Resulting concentrations were compared to New Jersey Soil Cleanup Criteria (NJDEP 1999; Table 3-15). The NJDEP criteria are human health based with categories addressing residential and non-residential settings, and impacts to groundwater. For our comparison, we reported the most conservative of the three criteria.

Table 3-15. Metals and semivolatiles (Group II) concentrations measured in sediments of Beltzville Reservoir in 2001.					
Parameter	Units	Method Detection Limit	BZ-6	USACE Screening Level	References
CONVENTIONALS					
Percent Solids	%	0.1	20.4		
METALS					
Aluminum	mg/kg	97.6	17,674.40		
Antimony	mg/kg	2	ND	14	NJDEP 1999
Arsenic	mg/kg	4.9	ND	20	NJDEP 1999
Barium	mg/kg	0.5	161.1	700	NJDEP 1996
Beryllium	mg/kg	0.5	1.1	2	NJDEP 1999
Cadmium	mg/kg	0.5	12.3	39	NJDEP 1999
Calcium	mg/kg	2	1513.6		
Chromium	mg/kg	0.5	26.1	33	MacDonald 1992
Cobalt	mg/kg	2	26.9		
Copper	mg/kg	0.5	32.8	600	NJDEP 1999
Iron	mg/kg	24.4	35,251.20		
Lead	mg/kg	2	105.5	400	NJDEP 1999
Magnesium	mg/kg	2	2,216.60		
Manganese	mg/kg	0.5	907.2		
Mercury	mg/kg	0.1	0.14	14	NJDEP 1999
Nickel	mg/kg	0.5	35.1	250	NJDEP 1999
Potassium	mg/kg	2	1152.3		
Selenium	mg/kg	4.9	ND	63	NJDEP 1999
Sodium	mg/kg	2	130.8		
Vanadium	mg/kg	2	45.7	370	NJDEP 1999
Zinc	mg/kg	0.5	682.6	1500	NJDEP 1999

Table 3-15. (Continued)					
Parameter	Units	Method Detection Limit	BZ-6	USACE Screening Level	References
SVOC (mg/kg)					
2,4,5-Trichlorophenol	mg/kg	0.490	ND	5,600	NJDEP 1999
2,4,6-Trichlorophenol	mg/kg	0.490	ND	62	NJDEP 1999
2,4-Dichlorophenol	mg/kg	0.490	ND	170	NJDEP 1999
2,4-Dimethylphenol	mg/kg	0.490	ND	10	NJDEP 1999
2,4-Dinitrophenol	mg/kg	0.490	ND	10	NJDEP 1999
2-Chlorophenol	mg/kg	0.490	ND	280	NJDEP 1999
2-Methylphenol	mg/kg	0.490	ND	2,800	NJDEP 1999
2-Nitrophenol	mg/kg	0.490	ND		
3-Methylphenol	mg/kg	0.490	ND		
4,6-Dinitro-2-methylphenol	mg/kg	0.490	ND		
4-Chloro-3-methylphenol	mg/kg	0.490	ND	10,000	NJDEP 1999
4-Methylphenol	mg/kg	0.490	ND	2,800	NJDEP 1999
4-Nitrophenol	mg/kg	0.490	ND		
Benzoic acid	mg/kg	0.490	ND		
Benzyl alcohol	mg/kg	0.490	ND	10,000	NJDEP 1999
Pentachlorophenol	mg/kg	0.490	ND	6	NJDEP 1999
Phenol	mg/kg	0.490	ND	10,000	NJDEP 1999

A total of 21 metals were analyzed in Beltzville Reservoir sediments (Table 3-15). None of compounds analyzed exceeded the screening level.

A total of 17 semivolatile organics (SVOC) were analyzed in Beltzville Reservoir sediments (Table 3-15). None of the compounds were detected in the sediment sample from station BZ-6 (Table 3-15).

3.6 DRINKING WATER

Drinking water from the public water fountain located in the overlook building of Beltzville Reservoir was monitored for compliance with PADEP water quality standards for primary and secondary contaminants, and quarterly monitored inorganic nitrogen (nitrate and nitrite) and coliform bacteria contaminants during 2001. Drinking water samples were analyzed in duplicate, comprising initial and confirmation samples. For matters of reporting, only if the result of the initial sample was not in compliance with water quality standards, the result of the confirmation sample was also reported.

3.6.1 Primary and Secondary Contaminants

Beltzville Reservoir drinking water was in compliance with PADEP water quality standards for all the primary and secondary contaminants (Table 3-16). As part of drinking water compliance monitoring, Safe Drinking Water Act (SDWA) form 4 for the reporting of results of primary and secondary drinking water contaminants were submitted to appropriate state environmental agencies.

Table 3-16. Concentrations of primary and secondary contaminants in drinking water at Beltzville Reservoir in 2001. Shaded values indicate results that exceeded Pennsylvania State drinking water standards; in these instances the result of a second sample is also reported.

Parameter	Sampling Date	PADEP Regulatory Level	Detection Limits	EPA Method
	14 June			
Aluminum	ND	0.2	0.02	200.7
Antimony	ND	0.006	0.05	200.7
Arsenic	ND	0.05	0.05	200.7
Barium	ND	2.0	0.005	200.7
Cadmium	ND	0.005	0.005	200.7
Chromium	ND	0.1	0.005	200.7
Copper	0.589	1.3	0.005	200.7
Iron	0.143	0.3	0.005	200.7
Lead	0.004	0.015	0.001	200.8
Magnesium	2.02	NL	0.02	200.7
Manganese	0.042	0.05	0.005	200.7
Mercury	ND	0.002	0.0002	245.1
Nickel	0.006	0.1	0.005	200.7
Selenium	ND	0.05	0.05	200.7
Silver	ND	0.1	0.005	200.7
Sodium	3.49	NL	0.02	200.7
Thallium	ND	0.002	0.05	200.7
Zinc	2.476	5.0	0.005	200.7
Chloride	2	250	1	325.3
Cyanide	ND	0.2	0.007	SM 4500CN-C&E
Fluoride	ND	2.0	0.1	SM 4500F-C
Foaming Agents	ND	0.5	0.05	SM 5540C
PH	7.14	6.5-8.5	+/-0.01	150.1
Sulfate	11	250.0	5	375.4
Total Dissolved Solids	34	500.0	10	SM 2540C

All results, criteria and detection limits are expressed in mg/L except pH which is expressed in positive/negative
 ND – Not Detected

3.6.2 Inorganic Nitrogen and Coliform bacteria

Beltzville Reservoir drinking water was in compliance with PADEP criteria for nitrate and nitrite (Table 3-17). Concentrations of nitrate and nitrite were not detected throughout the monitoring period. On 14 June during a scheduled sampling event, total coliform was present in the drinking water sample. A test collected on 21 June also detected coliform bacteria. To further investigate the source of bacteria four additional samples were collected on 26 June (Table 3-18). Water was collected from the observation fountain, the office fountain, the office sink, and the ranger's house faucet and tested for total coliform and fecal coliform. Coliform was present in the samples collected from the office sink. An additional sample collected on 5 July indicated that total coliform present. Subsequently, in late July, the USACE personnel at the Beltzville Reservoir office added a chlorine system to the water supply. On 7 August and there was no coliform bacteria contamination present. Additionally, coliform was absent on the scheduled 24 October sampling event. Following laboratory testing, drinking water monitoring results were recorded on Safe Drinking Water Act (SDWA-S and SDWA-4) forms and submitted to the appropriate state environmental agencies.

3.6.3 Historical Drinking Water Quality

Drinking water quality has been monitored at Blue Marsh Reservoir over the past 20 years. Versar (1996) compiled the results from all of the previous years into a single database to facilitate water quality comparisons. Historical data from drinking water quality parameters were compared to their respective PADEP standards. Of 26 parameters summarized, 7 had incidences of noncompliance with drinking water standards from 1983 to present (Table 3-18). Lead, pH, and corrosivity were most often not in compliance with PADEP criteria.

Table 3-17. Concentrations of nitrate and nitrite, and results of coliform bacteria monitoring of drinking water sampled from the public water fountain located in the overlook building at Beltzville Reservoir during 2001

Parameter	Sampling Dates					PADEP Regulatory Level	Detection Limits	Method
	14 June	21 June	5 July	7 August	24 October			
Nitrate as N (mg/L)	ND			ND	ND	10.0	0.5	SM4500
Nitrite as N (mg/L)	ND			ND	ND	1.0	0.5	SM4500
E. coli (CFU)	Absence	Absence	Absence	Absence	Absence	Presence	1	SM 9223
Total Coliform (CFU)	Presence	Presence	Presence	Absence	Absence	Presence	1	SM 9223
CFU – colony forming units ND – Not Detected								

Table 3-18. Results of coliform bacteria monitoring of drinking water sampled on 26 June at various places at the Beltzville Reservoir after the testing on 21 June. (Detection limit 10-clns/100-mls)

	Observation Fountain	Office Fountain	Office Sink	Rangers House Faucet
Total Coliform (10-clns/100-mls)	ND	ND	40	ND
Fecal Coliform (10-clns/100-mls)	ND	ND	10	ND
ND – Not Detected				

